


A Complete Ecbolic & Uterine Cleansing Tonic

Retention of fetal membranes, or retained placenta, usually is defined as failure to expel fetal membranes within 24 hrs after parturition. Normally, expulsion occurs within 3-8 hr after calf delivery. The incidence in healthy dairy cows is 5% - 15%, whereas the incidence in beef cows is lower. Retained placenta is most commonly associated with dystocia, milk fever (metabolic diseases) and twin births. Cows with retained fetal membranes are at increased risk of developing metritis, ketosis, mastitis, and even abortion in a subsequent pregnancy.



FytoBolic Forte is a synergistic combination of extracts of specific Phyto-active herbs, chelated minerals that stimulate the contraction of the uterus, post calving care of the uterus, relief in pain, inflammation, improve the milk production & prevent postpartum disorders.

CATALYST

Beyond Innovations

Composition:

Each 5 ml contains:

Zinc chelate	25 mg	Asparagus racemosus	40 mg	Hibiscus rosa sinensis	10 mg
Copper chelate	15 mg	Withania somnifera	40 mg	Acacia catechu	10 mg
Iron chelate	30 mg	Symplocos racemoses	20 mg	Holarrhena antidysentrica	5 mg
Cobalt chelate	05 mg	Terminalia chebula	20 mg	Embelia ribes	5 mg
Jivanti	10 mg	Terminalia bellerica	20 mg	Tinospora cordifolia	5 mg
Satavari	10 mg	Phyllanthus emblica	20 mg		
Peganum harmala	15 mg	Phyllanthus niruri	15 mg		
Citrllus colocynthis	250 mg	Ceiba pentandra	10 mg		
Aristolochia indica	250 mg	Glycyrrhiza glabra	15 mg		
Cedrus deodara	150 mg	Azadirachta indica	15 mg		
Dashmol	100 mg	Sida cordifolia	10 mg		
Syzygium cuminii	40 mg	Hemidesmus indicus	10 mg		

Benefits

- Aid in clearance of uterine discharge
- Hastens uterine involution
- · Restores milk production associated with uterine disorders
- Reduces postpartum hemorrhage
- Reduces the incidence of anestrus & repeat breeding
- · Helps as an adjunct during Metritis, Endometritis & Pyometra treatment

Indications:

- Facilitates expulsion of retained placenta
- For timely uterine involution
- To Manage pain & inflammation

Recommended Usage:

- Cows, Buffaloes and Mares: 100 ml
- Ewes and Does: 50 ml
- To be administered soon after calving
- In case of retained placenta, administer

double dose (200 ml for large animals and 100 ml for small animals) twice on first day followed by single dose twice daily for 3-5 days or as directed by the veterinarian.

out of smoke). When applying this product to water bodies, avoid solid particles from falling on nearby ground where birds and feed may be present.


Presentation: 500 ml poly can

Catalyst LifeSciences Pvt. Ltd.

An ISO 9001:2015, ISO 14001:2015, HACCP Compliance Certified Company

Corporate Office: Plot No.69, 1st Floor, Street No.1, Sector - 22, Noble Farms/Enclave, Gurugram - 122015, Haryana, India Customer care #: +91 892 988 1235; Email: support@catalystlifesciences.in

For more information please visit: www.catalystlifesciences.in DISPOSAL OF BAD AND OUTER PACKAGE: Do not contaminate water, food or feed by storage and disposal. Completely empty container into application equipment. Triple rinse pail and offer for recycling or reconditioning. Containers can also be disposed of in a sanitary landfill or by incineration, or in accordance with guidance from your local waste regulation authority, such as by burning (if burned, stay



Free From Hormones & Antibiotics

Animal Feed Supplement

Not For Medicinal use

Not For Human Use

